

4. Crawford, J. C. Descriptions of new Hymenoptera, No. 9. Proc. U. S. Nat. Mus. 43:577-586. 1915.
5. Crawford, J. C. New North American bees. Proc. Ent. Soc. Wash. 34:69-78. 1932.
6. Custer, C. P. Habits of *Perdita zebrata* with description of the larva. Can. Ent. 61:49-51. 1929.
7. Friese, H. Die Insekten Mitteleuropas insbesondere Deutschlands. Band. 1. Die Bienen, Wespen, Grab-und Goldwespen. Stuttgart, 1926.
8. Michener, Chas. D. Comparative external morphology, phylogeny, and a classification of the bees. Bull. Amer. Mus. Nat. Hist. 82:151-326. 246 figs. 1944.
9. Stevens, O. A. The panurgine bees of North Dakota and a new *Epeolus*. Can. Ent. 51:205-210. 1919.
10. Timberlake, P. H. Bees of the genus *Perdita* in the J. C. Bridwell collection. Proc. Haw. Ent. Soc. 7:151-161. 1928.

CORRECTION

Two errors in the article on *Andrena*, Vol. XII, No. 1, were: Page 19, next to last line in No. 25, should read "series is" in place of "species are".

Page 22, the record for Slope County under No. 55, should have been under No. 54, *Andrena haynesi*.

NORTH DAKOTA WINTER RYE

North Dakota's winter rye acreage of 279,000 acres seeded the fall of 1949 was about six per cent over the previous year but still far below the 679,000 acres for the 10-year average, says C. J. Heltemes, USDA statistician at Fargo. He believes dry weather in the fall cut winter rye plantings. While North Dakota ranked first in rye production in the nation in 1949, our 1949 fall seeding was third, being exceeded by South Dakota and Nebraska, where fall planting was increased 40 and 12 per cent respectively.

1949 PIG REPORT

Eleven per cent more pigs were raised in North Dakota in 1949 than in 1948, according to the USDA bureau of agricultural economics. The combined spring and fall crops amounted to 855,000 head, compared with 767,000 in 1948 but still significantly below the 10-year average of 1,044,000. Iowa led the nation in 1949 with over 19 million head, followed by over 10 million in Illinois.