

Hard Red Spring Wheat and Durum Wheat Production Guide (continued)

A-1050, May 1998

Weed Control (continued)

Post Applied Herbicides Continued

	L a n c e l e a f S a g e	M a l l o w, V e n i c e	M a r s h e l d e r	M u s t a r d, W i l d	N i g h t s h a d e, B l a c k	P i g w e e d, R e d r o o t	P r i c k l y L e t t u c e	R a g w e e d, C o m m o n	S m a r t w e e d, A n n u a l	S u n f l o w e r	T h i s t l e, R u s s i a n	T h i s t l e, C a n a d a	H e r b i c i d e P e r s i s t e n c e
POST APPLIED HERBICIDES													
Achieve	N	N	N	N	N	N	N	N	N	N	N	N	N
Ally + 2,4-D	F-G	G-E	G-E	E	F-G	E	E	E	F	G-E	E ²	G	O
Amber + 2,4-D	F-G	—	E	E	F-G	E	E	E	F	E	E ²	F-G	O

Assert	N	N	N	E	—	P	N	N	P	N	P-F	N	S
Avenge	N	N	N	N	N	N	N	N	N	N	N	N	N
Banvel/SGF/Clarity	P-F	F	E	G	E	G	G-E	E	E	G-E	G	F-G	S
Banvel/SGF + MCPA	G-E	G	G-E	E	E	G	E	E	E	E	G	F	S
Bronate/others	E	G-E	E	E	E	G	E	E	G	E	E	F	N
Bromoxynil	E	G-E	E	F-G	E	F-G	E	E	G-E	G-E	E	F	N
Canvas + 2,4-D	F-G	G-E	E	E	F-G	E	E	E	F-G	G-E	E ²	G	O
Cheyenne	F-G	G	E	E	F-G	E	E	E	E	E	G-E ²	P	N
Curtail/M	F-G	G	E	E	F	G	E	E	E	E	G	G-E	S
Dakota	F-G	F	G	E	F	P	G-E	G	G	G	P	P	N
Express + 2,4-D	F-G	—	E	E	F-G	G	G-E	G	G	F-G	E ²	G	N
Finesse + 2,4-D	F-G	E	E	E	F-G	E	E	E	E	E	E ²	G	O
Fusilade DX	N	N	N	N	N	N	N	N	N	N	N	N	N
Gramoxone Extra	E	G	G	E	G-E	E	E	G-E	E	E	E	P	N
Harmony Extra + 2,4-D	F-G	—	E	E	F-G	E	E	E	G	G-E	E ²	G	N
Hoelon	N	N	N	N	N	N	N	N	N	N	N	N	N
MCPA	G-E	F	G	E	F	F	F	G	F	G	P	F-G	N
Peak + 2,4-D	F-G	—	—	E	—	E	E	E	—	E	E ²	F-G	O
Puma	N	N	N	N	N	N	N	N	N	N	N	N	N
Roundup Ultra/RT/Glyphos	E	E	G-E	G-E	F-G	G-E	E	E	E	G	F-G	G	N
Stampede EDF + MCPAe	N	—	F	E	—	E	—	P	F	G	F	F	N
Stinger	F	G	E	P	F-G	P	E	G-E	G-E	G-E	P-F	G-E	S
Tiller	G	F-G	F-G	E	F	F	E	F	P	G-E	F	F	N
Tordon 22K + 2,4-D	—	G-E	E	E	G-E	E	E	G-E	G	E	G-E	F-G	O
Touchdown	E	E	G-E	G-E	F-G	G-E	E	E	E	G	F-G	G	N
2,4-D	P-F	G-E	E	E	P-F	G	E	G-E	F-G	E	G	F-G	N

¹ Ally, Amber, Express, Finesse, and Harmony Extra tank-mixed with 2,4-D and Banvel/SGF will provide excellent control of most broadleaf weeds and reduce risk of developing ALS resistant kochia.

² Herbicides will not control resistant biotypes.

³ Tordon + 2,4-D at 1 to 1.5 fl oz/A + 2,4-D at 0.5 to 0.75 pt/A

[[CONTINUE](#)] [[INDEX](#)]

County Commissions, North Dakota State University and U.S. Department of Agriculture cooperating. North Dakota State University does not discriminate on the basis of race, color, national origin, religion, sex, gender identity, disability, age, status as a U.S. veteran, sexual orientation, marital status, or public assistance status. Direct inquiries to the Vice President for Equity, Diversity and Global Outreach, 205 Old Main, (701) 231-7708. This publication will be made available in alternative formats for people with disabilities upon request, 701 231-7881.

INFORMATION ACADEMICS RESEARCH EXTENSION PUBLICATIONS CALENDAR WEATHER DIRECTORY

[Information for Prospective Students](#)

NDSU is an equal opportunity institution

This information may be copied for noncommercial, educational purposes in its entirety with no changes.

Requests to use any portion of the document should be sent to NDSU.permission@ndsu.edu.

North Dakota State University Agriculture and University Extension

NDSU Dept. 7070, P.O. Box 6050, Fargo, ND 58108-6050