

Grasses (Forage) SEED TREATMENT

Chemical	Application	Dosage ¹	Control ² of Seedling Blights ³	Remarks
Fludioxonil Maxim 4 FS, 40.3%	Slurry	0.08-0.16 fl oz/cwt	X	For seed-borne and soil-borne fungi.
Mefenoxam Apron XL LS, 32.3%	Slurry or mist	0.32-0.64 fl oz/cwt	X	Apron XL LS controls only Pythium. For both commercial and on-farm use.
Metalaxyl Acquire, 28.35% Allegiance FI 28.35% Sebring 318 FS, 28.35% Allegiance Dry Seed Protectant, 12.5% Dyna-Shield, 28.35%	Mist or slurry	0.75 fl oz/cwt	X	Acquire and Allegiance controls only Pythium.
	Drill box	3-4 oz/cwt	X	Allegiance Dry Seed Protectant is for drill box application to seed not previously treated with Apron; thorough mixing of fungicide and seed is essential for good control.
	Slurry	0.75 fl oz/cwt	X	
Thiram 42-S Thiram, 42% Thiram 50WP Dyed, 50%	Liquid or slurry	8 fl oz/cwt	X	
	Drill box or slurry	8 oz/cwt	X	

¹Dosage = amount of formulated product to apply.

²X = product labeled for crop and disease; Blank = product not labeled for specific disease.

³Seedling blights due to various fungal infections of seed.