

2009 North Dakota FIELD CROP FUNGICIDE GUIDE

Compiled by
Marcia P. McMullen and Samuel G. Markell
Extension Plant Pathologists

in cooperation with

the faculty of the
Plant Pathology Department
North Dakota State University

TABLE OF CONTENTS

Disclaimer	1
Label Precautions and Restrictions	2
-Field re-entry, handling and loading precautions; Replant restrictions; Dosages; Restricted use fungicides	
Fungicide Formulations	2
-Wettable powders (WP); Water Soluble Pouch (WSP); Dusts (D); Granules (G); Emulsifiable concentrates (EC); Flowables.	
Mode of Action of fungicides	3
Toxicity of Fungicides- Effects of Chemicals on Humans; First aid, Toxicity ratings of pesticides	3
Protecting Ground Water, Handling chemicals	4
Fungicide Labels	4
Seed Treatment: Cereals; Chickpeas; Dry Edible Beans and Soybeans; Flax; Potatoes; Safflower; Sunflower	4
Application of Seed Treatment	5
Field Crop Foliar Sprays	6
Resistance to fungicides	6
Fungicide Resistance Management	7
Fungicide Groups	7
Section 18 Emergency Uses Requested for 2009	8
Tables :	
-Alfalfa-Clover-Small Seeded Legumes Seed Treatment	9
-Barley Seed Treatments	10
-Barley Foliar Sprays	15
-Canola Seed Treatments	19
-Canola Foliar Sprays	20
-Chickpea Seed Treatments	21
-Chickpea Foliar Sprays	22
-Corn and Sorghum Seed Treatments	23
-Corn and Sorghum Foliar Sprays	25
-Crambe Seed Treatments	26
-Dry Edible Bean Seed Treatments	26
-Dry Edible Bean Biological Seed Treatment	28
-Dry Edible Bean Soil Application	28
-Dry Edible Bean Foliar Sprays	29
-Flax Seed Treatments	32
-Grasses (Forage) Seed Treatments	33
-Lentil Seed Treatments	34
-Lentil Foliar Sprays	35
-Oat Seed Treatments	10
-Oat Foliar Sprays	15
-Pea Seed Treatments	36
-Pea Biological Seed Treatment	37
-Pea Foliar Sprays	37
-Potato Seed Treatments	38
-Potato Soil Application	39
-Potato Foliar Sprays	40
-Rye Seed Treatments	10
-Rye Foliar Sprays	15
-Safflower Seed Treatments	47
-Safflower Foliar Sprays	47
-Soybean Seed Treatments	48
-Soybean Biological Seed Treatments	49
-Soybean Soil Application	50
-Soybean Foliar Sprays	51
-Sugarbeet Seed Treatment	53
-Sugarbeet Soil Application	54
-Sugarbeet Foliar Sprays	55
-Sunflower Seed Treatment	58
-Sunflower Foliar Sprays	59
-Wheat Seed Treatment	10
-Wheat Foliar Sprays	15
-Soil Applied Biological Fungicide	59
Distributor List	60

This fungicide guide is based on the latest information available from the North Dakota Agricultural Experiment Station, U.S. Department of Agriculture, U.S. Environmental Protection Agency (EPA) and the agricultural chemical industry. The information conformed to federal and state regulations at the time of printing. The user should determine that the intended use is consistent with label directions. ***Designation that a product is labeled for control of a crop disease does not imply endorsement by the authors of use of that product or the degree of efficacy of that product for that use.***

Always follow the label directions. See individual fungicide labels for important information on:

- Safety recommendations and worker protection requirements
- Guidelines for ground, irrigation, or aerial application
- Mixing procedures and tank mixes allowed
- Rotational and grazing restrictions
- Resistance management statements

LABEL PRECAUTIONS, RESTRICTIONS

Field re-entry, handling and loading precautions

Most fungicide labels state that workers either should not enter a sprayed field until the sprays have dried or should not enter for 24 hours unless they wear appropriate protective clothing. Information on use of protective clothing during mixing and loading also is given on the label. See the label for details.

Replant restrictions

Labels for all formulations of Ridomil have restrictions on what crops can be planted in less than a year following application of the product. These restrictions may vary somewhat depending on the formulation. Check these and all other labels **before** application to determine if replant restrictions will cause problems when determining what crop to plant next season.

Dosages

All dosages given in this guide are stated as the amount of formulated product (lb., oz., fl. oz., quarts) to use.

Restricted use fungicides

Fungicides containing triphenyltin hydroxide are restricted-use fungicides. These include products such as Super Tin, Agri Tin and Super Tin 4L. These are designated as RUP and Restricted-use Pesticide in the tables.

Disclaimer

The information given herein is for educational purposes only. North Dakota State University does not endorse commercial products or companies, even though reference may be made to trade names, trademarks or service names. **Omission of labeled products is possible if information about the product was not available at the time of printing or if it had questionable efficacy. Products not normally available in North Dakota are omitted from the guide. Seed treatment chemicals that are primarily insecticides with subminimal amounts of fungicide are also omitted.**

The plant pathology faculty at North Dakota State University assumes no responsibility for property damage, personal injury or other loss due to the use of fungicides listed in this publication, as they have no control over the use or misuse of these products.

FUNGICIDE FORMULATIONS

Most fungicides are solids that are not soluble in water. To use them, they must be made into a formulation (preparation). Some of the more common formulations are listed below. The common abbreviation for each formulation is given in parentheses following the name.

Wettable powders (WP)

Many fungicide are wettable powders consisting of solid fungicide and a wetting agent. When mixed with water, they form a suspension. Many of these suspensions settle out quickly, so an agitator is needed in the spray tank to keep the particles in suspension.

Water Soluble Pouch (WSP)

Some fungicides are available in water-soluble pouch containers. These pouches dissolve in the mixing tank and release the fungicide. This reduces the exposure of mixer and loader personnel to dust from the fungicide.

Dusts (D)

Dusts are powders that are mixed with inert ingredients to form a product with a low percent of active material. These are used around the home garden, and a few formulations are used in commercial applications.

Granules (G)

The active ingredient is incorporated into small granules of inert material such as clay. Granules are incorporated into the soil.

Emulsifiable concentrates (EC)

A fungicide that is insoluble in water is dissolved in an organic solvent. An emulsifying agent is incorporated in the formulation, so an emulsion is formed when the product is mixed with water. An emulsion is a suspension of very tiny drops of the solvent/fungicide in the water. It usually has a milky appearance (milk itself is an emulsion of fats in water).

Flowables (F)

Flowables are insoluble fungicides ground into a very fine product, usually by a wet grinding process. These particles are nearly colloidal in size and are suspended in water to form a thick liquid. They remain suspended in water for relatively long periods of time, but should be agitated before use. They are dust-free, easy to mix, remain in suspension longer than wettable powders, and also may resist washing off the plant better than the wettable powders. Examples of flowables include Champ Flowable, Kocide 4.5 LF, Vitavax 200, and Dithane F-45. They need to be protected from freezing.

Dry flowable (DF)

See dispersible granules. (Next page)

Dispersible granules (DG)

Dispersible granules also are called dry flowable formulations. They are small-sized granules that pour from a container like a liquid, but do not stick to the sides of the container and do not need to be protected from freezing. They are virtually dust-free and disperse readily in water to form a suspension. Examples include Bravo Ultrex DG, Dithane DF, Rainshield NT, Manzate 75 DF, and Penncozeb DF.

Fumigants

Fumigants are liquids that turn into a gas after application. They generally are used for soil fumigation.

MODE OF ACTION OF FUNGICIDES

The action of most fungicides takes place outside the host and is called "*protection*." A fungicide that acts outside the host is called a "protectant fungicide." Most older fungicides sprayed on leaves and fruit are of this type. "*Therapy*" is chemical action inside the host. For example, fungicides are locally systemic and move into the plant at the site of deposition. Several triazole fungicides have several days of therapeutic action against wheat leaf rust and also reduce the production of viable spores; that is, spores capable of growing.

Most protectant fungicides are relatively stable by themselves. Generally, they are relatively insoluble in water and resist removal or chemical change by water, yet must be toxic to fungi. Often a chemical change is brought about by the fungus, the host, or the environment before toxicity occurs. Toxicity simply means the ability to damage the fungus cells.

Fungicides may act to produce a toxic reaction in the fungus in several different ways. (1) Some may inhibit (slow down or stop) cell wall formation. (2) Some affect the permeability of the cell wall, causing a leaking of nutrient materials from the cell. (3) Some fungicides may combine with essential metals in a way that they become unavailable for normal cell functions, including the functioning of essential enzymes. (4) Other fungicides may inhibit respiration, or nuclear division, or may break dormancy of spores.

Some fungicides also may be toxic to plants if applied at rates too high, or if applied under unfavorable environmental conditions. This is called *phytotoxicity*. Formulations of maneb + zinc are less phytotoxic to many vegetables than formulations that contain only maneb. Sometimes the method of formulation may make a fungicide less phytotoxic.

TOXICITY OF FUNGICIDES

Effects of Chemicals on Humans

Fungicides have various levels of toxicity to humans. Human exposure (skin, eye, internal) to fungicides can

result in mild to severe reaction. Due to high levels of toxicity, some fungicides are restricted use only.

Symptoms associated with chemical poisoning are listed below. All symptoms are not associated with every pesticide. Some of these symptoms are described below, but it is always wise to consult a physician. Avoid diagnosing the effects on yourself or others.

- Eyes watering excessively
- Stomach cramps
- Dizziness
- Vomiting
- Excessive sweating
- Pupils of the eye reduced in size
- Rapid heart beat
- Muscle tremors or convulsions
- Extreme nervousness
- Mental confusion, lack of coordination
- Uncontrolled drooling or watering at the mouth
- Severe burns of the skin
- Loss of ability to use muscles
- Difficulty in breathing
- Unconsciousness

First aid

The following list should be considered:

- Stop exposure
- Call a physician
- Remove contaminated and restrictive clothing
- Drench contaminated area with water, flush repeatedly
- Provide fresh air, but prevent chilling and overheating
- Avoid giving alcohol
- Provide milk for patient to drink
- Antidote - to be administered only by a physician.

NORTH DAKOTA POISON CONTROL CENTER
TOLL FREE: (800) 732-2200

Toxicity ratings of pesticides

Pesticides generally are categorized according to acute **oral toxicity** (the toxicity when taken by mouth), but because users may absorb a significant quantity of the pesticide through their skin, **dermal toxicity** (toxicity when absorbed through the skin) is of equal or greater practical importance.

LD₅₀ values generally show relative toxicities among the chemicals and are not truly representative of effects on humans, especially since they usually are obtained on rats. Actual toxicities do not constitute the only hazards associated with exposure to the chemicals. For instance, a chemical with low toxicity may be hazardous due to concentration, high volatility, careless use, or effects of long-term exposure.

LD₅₀ depends upon body weight. Thus, a given amount of chemical would have greater effect on a child than on an adult. LD₅₀ also is proportional to the percent of active ingredient. A material only 50 percent active requires twice as much to produce a toxic effect as 100 percent pure material.

The lower the LD₅₀ value, the greater the toxicity. A common standard for comparison is aspirin, which has an LD₅₀ of 1,200 mg/kg and is considered slightly toxic.

The following table illustrates the various toxicity classes:

Oral Toxicity		Dermal (Skin) Toxicity	
LD ₅₀ -mg/kg	Toxicity Class	LD ₅₀ -mg/kg	Toxicity Class
1-50	High	1-200	Severe
50-500	Moderate	200-2,000	Moderate
500-5,000	Low	2,000-20,000	Mild
Over 5,000	Very Low	Over 20,000	Very Mild

Information on the LD₅₀ of a specific fungicide, and other toxicology information, is available on the MSDS (Material Safety Data Sheet) for each product. These generally may be found at <http://www.cdms.net/>

Protecting Groundwater

Pesticides differ in their persistence and mobility in soil. Those which are highly persistent or highly mobile are more liable to contaminate groundwater than those that are not. Areas of the state where groundwater is most at risk are areas with coarse-textured soils, are low in organic matter and have a high water table. Most fungicides are relatively immobile, especially in clay soils with high organic matter because they are adsorbed on clay particles or on the organic matter.

A few fungicides are somewhat mobile. Take care in the use of these fungicides, particularly the application of these products through a sprinkler irrigation system in high-risk areas. Risks may be reduced by minimizing the amount of water used for application through a sprinkler system, more use of ground or aerial application instead of application through the sprinkler system, and use of a different fungicide that is less mobile.

The persistence and mobility of fungicides commonly used in North Dakota may be found in NDSU Extension Service publication EB-49 "Persistence and Mobility of Pesticides in Soil and Water."

Handling chemicals

Avoid splashing and spilling. Wear a mask especially when handling dusts or powders. Some chemicals, when combined, have increased toxicity (potentiation).

Rinse containers several times after using chemicals. Pour rinsate into spray tank when using the same chemical. Dispose of containers as indicated in next

section. Keep a record of plant disease control chemicals used and methods of handling.

Fungicide Labels

Fungicides are named according to their chemical composition or the *chemical name*. An example of a chemical name is a coordination production of zinc ion and manganese ethylene bisdithiocarbamate; the chemical names are required on the label. Since chemical names are frequently long, *common names* frequently are used; for example, the common name for the above chemical is mancozeb. Manufacturers use *trade names* to identify their specific products. For example, there are various trade names for mancozeb such as Dithane, Manzate, and Penncozeb.

In addition to the names on labels, various other required label information includes: precautions in handling, antidotes or telephone contacts to use in case of accidental poisoning, recommendations for use, materials contained in the package and their percentages, the manufacturer's or distributor's name and address, and the EPA registration number.

Some fungicides are made up in various formulations for different uses or methods of application such as: wettable powders, dusts, emulsifiable concentrates, granules, flowables, dispersible granules, or solutions. The nature of the chemical sometimes restricts it to one or a few of these formulations.

Note: Although maneb and mancozeb are very similar products, there are some differences in registration. Be sure to consult and follow label directions.

Seed Treatment

Cereals

Fungicidal seed treatment helps protect the seed from rotting and the emerging seedlings from damping off and seedling blight. These are caused by soil-borne pathogens. When seeds germinate under favorable soil conditions, there is less danger of seed and seedling attack from soil-borne pathogens unless seed is of poor quality. Treatment of seed with a protectant fungicide may help protect against soil-borne pathogens and thus help stand establishment when seeds are germinating under unfavorable conditions, such as cold wet weather. Many products are available for protection against seedling blight.

Treating seeds with a fungicide also helps protect them from diseases that are seed-borne. These include the covered smuts, bunt, scab, black point, and black semi-loose smut of barley, and loose smuts of wheat, barley and oats. Loose smuts of wheat and barley are internally seed-borne. Loose smut of oats is seed-borne as spores under the hulls. These smuts cannot be controlled by conventional protectant seed treatment fungicides, but are controlled by systemic seed treatment products. The

embryo test can be used by the North Dakota State Seed Department to determine if loose smut is present in barley seed. This test cannot be used for the loose smuts of oats or wheat or black semi-loose smut of barley. All current barley varieties are susceptible to loose smut. An embryo test is recommended for barley seed; if infection is 2 percent or greater, seed treatment of barley with carboxin or triadimenol is advised.

Common (*Bipolaris*, *Helminthosporium* or *Cochliobolus*) root rot of wheat and barley is a chronic problem in North Dakota, causing average yield losses of 5 to 11 percent, with much greater losses in some fields in certain years. Several seed treatment products are labeled for suppression of common root rot.

Chickpeas

Treating chickpea seed to protect against *Pythium* is essential for good emergence. A seed treatment to protect against seed-borne *Ascochyta* is important as this is a common and serious disease.

Dry beans and soybeans

Treating seed may reduce seedling blight during weather that is unfavorable for emergence. Do not use streptomycin with Rhizobium inoculant. If using captan seed treatments, in-furrow inoculant is preferable, because inoculant does not survive well on captan-treated seed. Several products can be used to reduce the root rot potential, and many newer products have a broad spectrum of activity.

Flax

Treating flax seed with a fungicide helps protect against seed rot, damping off, and seedling blight. Seed treatment is especially important in cases where the seed coats are broken, allowing entry of pathogens. Seed from fields heavily infected with Pasm (*Septoria linicola*) may be susceptible to seedling blight and should be seed treated.

Potatoes

Treatment of cut seed-pieces helps protect the cut surface against seed-piece decay. Most seed treatments are fungicides that will protect against fungi such as *Pythium*, *Rhizoctonia* and *Fusarium*. Fungicides do not protect against bacteria such as *Erwinia* or *Clavibacter*. However, control of fungi indirectly helps control *Erwinia* bacteria because seed decay is greater in seed infected with fungi. The addition of streptomycin to fungicide has limited value because it will control only bacteria contaminating cut surfaces and may inhibit wound healing. Seed treatment will reduce or help control new infections but will not cure existing decay, prevent lenticel infection or prevent infection of roots and stolons away from the seed piece due to soil or environmental inoculum. Seed treatment is no substitute for using good sound healthy seed. Seed should be stored at 40F during the winter. In the spring, warm the seed to 60-70F for 1 1/2-2 weeks before planting. Do not handle the seed until it is warm. Plant the cut seed in warm (50-58F at planting depth) moist soil. If cut seed must be held, store in a well-ventilated area for suberization at 60-70F with a relative

humidity of 85 percent. Hold for 1 week, then lower the temperature to 50-60F.

Safflower

Safflower rust is both seed-borne and soil-borne. The most devastating phase of the disease is a seedling blight, and root and foot rot. Typical rust pustules develop later on the leaves. Seed-borne safflower rust is controlled by seed treatment.

Sunflower

Soil-borne downy mildew infections were controlled with metalaxyl or mefenoxam seed treatment in the past. The downy mildew fungus, however, has developed insensitivity to metalaxyl and mefenoxam in much of North Dakota, South Dakota, and Minnesota, so these fungicides are not effective. Several fungicides or fungicide-insecticide combinations have received state or federal labels for seed treatment of sunflower for seed rot and seedling blights.

Application of Seed Treatment

Seed may be treated commercially or it may be treated on the farm. Commercial seed treatment may use a slurry treater or various automatic seed treaters. The various automatic seed treaters differ considerably and so cannot be discussed here. Commercial seed treatment has become more common in recent years for many crops.

On-farm treatment may use various home-type or slurry mixers. Drill-box seed treatment is popular because no extra steps are required because the seed is treated in the drill-box at planting time. Good disease control depends on uniform fungicide coverage of the seed, but this is more difficult to accomplish in drill-box treatment because there's an inadequate means of mixing the seed and fungicide. For effective drill-box treatment, fill the box with one-third the quantity of seed and fungicide and mix carefully with a paddle; repeat with the next third, and then the final third. The paddle should not be used for any other purpose and should be stored in a safe place, out of reach of children and animals.

On-farm auger seed treatment methods are common. The fungicide is metered into the base of the auger used to fill the drill box. This method assures fairly good mixing and coverage.

All seed treatments have certain basic precautions. Use care in handling seed treatment products — many are irritating to the eyes, nose and skin. Treated seed usually is identified by the dye used in the chemical, and treated seed should not be fed to livestock or used for human food. Pesticide containers should be disposed of properly in a landfill or buried in an area where there is no surface drainage to nearby water ways. If seed treatment cannot be done outdoors, it should be done in a well-ventilated room. Commercial seed treaters should have an adequate air exhaust system for treatment rooms. Workers exposed to seed treatment chemicals for long periods of time should have an approved chemical mask. The filter should be changed frequently. Recommended rates of application

should be followed carefully as higher rates may injure the seed and lower rates may not give satisfactory disease control.

Forage legume seed should be treated well in advance of planting and inoculated with nitrogen fixing *Rhizobia* at planting time. If dry beans have been treated with streptomycin for control of externally borne blight bacteria, inoculating with *Rhizobia* is not available.

Field Crops Foliar Sprays

Foliar fungicides are used to control fungal disease organisms that attack the above-ground portions of plants. Fungicides are used to protect the potential yield and quality of a crop. Many fungicides protect foliage from infection; therefore, these fungicides must be on the foliage before the fungus spores germinate.

Several foliar fungicides act differently from the protectants described above. For example, benzimidazole fungicides thiabendazole and thiophanate methyl are absorbed by the plant and translocated up the plant by the conducting tissues. They are called systemic fungicides. They only move up the plant; they do not move down. Thus, to control white mold on dry beans, complete coverage of stems, lower leaves and blossoms is required. Spraying only the upper leaves is not satisfactory because the fungicide will not move down to the location where it is needed. Strobilurin and triazole fungicides are locally systemic, they have some upward mobility and translaminar movement and have some limited therapeutic action. Metalaxyl will move down from potato foliage into tubers in limited amounts to provide tuber protection against metalaxyl- sensitive strains of the late blight fungus and pink rot infection.

Spray control programs to prevent disease have been developed from data through years of research. Because each disease develops in a distinct manner, the decision to use a disease prevention program is based on weather conditions, disease development, potential yield of the crop, and the dollars returned to management with use of the fungicides.

Many fungicides are registered for application through a sprinkler irrigation system, as well as by a spray. If a fungicide can be applied through a sprinkler system (fungigation), this is noted under application.

Most fungicide labels contain information on field re-entry, handling and loading precautions. Most labels state that workers either should not enter a sprayed field until the sprays have dried or should not enter for 24 hours unless they wear appropriate protective clothing. Information on the use of protective clothing during mixing and loading also is given on the label. See the label for details.

Spraying

Spraying can be done with many different types of ground and air equipment. Getting good coverage is important: use at least 5 gal/A should be used for aerial application and higher gallonages are required for ground equipment.

Droplet size for aerial application should be 200-400 microns (1/64 to 1/128 inch) in diameter. Generally, if nozzles are pointed back, appropriate nozzles are used, and pressures do not exceed 30 or 35 psi, the correct droplet size will result. Application should be made with the boom 6 to 8 feet above the crop.

Some plant surfaces have a waxy or hairy coating on them, making good coverage difficult. The spray will collect in large erect droplets which then run off. Wheat and cabbage leaves are good examples. Frequently, using a wetting agent with wettable powders is necessary. Usually this is a spreader-sticker. Certain fungicides may work better with certain spreader-stickers than others. This type of information usually can be found on the label or in supplemental brochures. Spreader-stickers may be incorporated into some flowable formulations, so adding a spreader-sticker to the spray tank is not necessary. However, the label must be checked on each product recommendation for this use.

Resistance to fungicides

Fungi may develop tolerance or resistance to certain fungicides. The sugarbeet leafspot fungus (*Cercospora*) has developed resistance to the systemic benzimidazole fungicides (benomyl, thiabendazole, and thiophanate methyl) in the Red River Valley and southern Minnesota. These fungicides should be not used at all in the southern Red River Valley and no more than once a season in a tank mix with an unrelated fungicide in the northern Red River Valley.

Resistance to the benzimidazole fungicides thiabendazole (TBZ or Mertect) and thiophanate methyl (Topsin M) has recently developed in the potato *Fusarium* dry rot pathogen, *Fusarium sambucinum*, and the potato silver scurf pathogen, *Helminthosporium solani*. This resistance is common throughout the United States and Canada.

Resistance to iprodione has been reported from other parts of the country. Cross-resistance to the chemically related product vinclozolin is common when resistance to iprodione develops.

The potato late blight fungus developed resistance to metalaxyl in Europe. The A2 mating type of the late blight fungus, which is common in North Dakota and Minnesota, is resistant to metalaxyl and mefenoxam.

In ND, reduced sensitivity to strobilurin fungicides have been observed in populations of the early-blight fungus *Alternaria* sp. on potato and to the aschochyta pathogen on chickpeas.

Tolerance of the sugarbeet leafspot fungus to triphenyltin hydroxide was widespread in southern Minnesota and the southern Red River Valley in 1999, and common in the northern Red River Valley. Tin-tolerant strains have not survived well when an alternative fungicide was used.

In contrast, benzimidazole-resistant strains survive very well when alternative fungicides are used, and persist for a long time. The best way to combat resistance is to prevent or delay it by alternating the use of different classes of fungicides and by avoiding constant use of fungicides known to trigger development of resistant fungi. Using tank mixes of unrelated fungicides also is reported to retard the development of resistance.

Fungicide Resistance Management Statements

The following statements are recommendations from the Fungicide Resistance Action Committee (FRAC), whose website is: <http://www.frac.info/>

1. Methyl benzimidazole carbamates (MBC; Group 1) – High risk. Both mixtures and alternations with non-Group 1 fungicides are acceptable methods of preventing/managing resistance to Group 1 fungicides. For high risk pathogens, mixtures are preferred to alternations.

2. Dicarboximides (Group 2) – Medium to high risk. Minimize the selection pressure by minimizing the number of applications. As a guide, do not apply more than two to three per crop per season. Maintain regular prolonged times without exposure to Group 2 fungicides. When applying for *Botrytis* control: Restrict applications to those times when *Botrytis* infection pressure is high. Where *Botrytis* resistance is well established, use combinations to stabilize *Botrytis* control, but their application must follow the same rules as for Group 2 fungicides alone.

3. Sterol biosynthesis inhibitors (SBI; Groups 3, 5, 17, and 18) – Low to medium risk. Repeated applications of SBI fungicides alone should not be used on the same crop in one season against a high-risk pathogen in areas of high disease pressure for that particular pathogen. For crop/pathogen situations where repeated spray applications are made during the season, alternation or mixtures with an effective non cross-resistant fungicide are recommended. Where alternation or the use of mixtures is not feasible because of lack of effective or compatible non cross-resistant partner fungicides, then input of SBIs should be reserved for critical parts of the season or crop growth stage. If SBI performance should decline and sensitivity testing has confirmed the presence of less sensitive forms, SBIs should only be used in mixture or alternation with effective non cross-resistant partner fungicides. The introduction of the new classes of chemistry offers new opportunities for more effective resistance management. The use of different mode of actions should be maximized for the most effective resistance management strategies. Users must adhere to the manufacturers' recommendations. In many cases, reports of "resistance" have, on investigation, been attributed to cutting recommended rates of use, or to poor or miss-timed application. Fungicide input is only one aspect of crop management. Fungicide use does not replace the need for resistant crop varieties, good agronomic practice, plant hygiene/sanitation, etc.

4. Phenylamides (PA; Group 4) – High risk. The Group 4 fungicides should be used on a preventative and not curative or eradication basis. For foliar applications, Group 4 fungicides should be used in pre-packed mixtures with an unrelated effective partner and used in a sound management program. Where using residual partners, use between three-fourths and full recommended rates. The Group 4 fungicide dosage in the mixture depends on its intrinsic activity and is defined by the respective company. The Group 4 fungicides should not be used as soil treatments against airborne diseases. When solo formulations are made available for soil use, strategies that prevent any possibilities for foliar applications must be implemented. For seed treatment, mixtures rather than straight Group 4 fungicides should be used whenever possible. The number of Group 4 fungicide applications should be limited (two to four consecutive applications per crop and year). The application intervals should not exceed 14 days and may be shorter in cases of high disease pressure. If rates and application intervals are reduced, the total amount of the Group 4 fungicide used per season should not exceed that of the full rate, and the total exposure time should remain the same. The rate of the mixing partners should remain the same for both intervals. Group 4 fungicide sprays are recommended early season during the period of active vegetative growth of the crop. The grower should switch to non-Group 4 products not later than the normal standard application interval of the non-Group 4 product.

5. Quinone outside inhibitors (QoI; Group 11) – High risk. When using a Group 11 fungicide as a solo product, the number of applications should be no more than one-third of the total number of fungicide applications per season. In programs with tank mixes or pre-mixes of a Group 11 fungicide, applications should be no more than one-half of the total number of fungicide applications per season. In programs in which applications of Group 11 fungicides are made with both solo products and mixtures, the number of Group 11 fungicide-containing applications should be no more than 1/2 of the total number of fungicide applications per season.

Fungicide Groups

The Soil Application and foliar sprays tables in this guide have a numerical or letter designation (in parentheses) for each chemical component of the listed commercial fungicides. This number or letter code indicates the fungicide group to which the chemical belongs. The fungicide groups have been designated by the Fungicide Resistance Action Committee (FRAC). FRAC is a Specialist Technical Group of the Global Crop Protection Federation. The members of this committee are representatives from all major crop protection companies worldwide. The FRAC code of numbers and letters are widely accepted in the literature.

The purpose of FRAC is to prolong the effectiveness of fungicides liable to encounter resistance problems and to limit crop losses should resistance appear. If field resistance is known to one member of the Fungicide Group, it is possible that cross resistance to other

chemicals within that Group will be present. This Fungicide Guide is providing information on Fungicide Groups so that users are aware of potential resistance problems with continued use of chemicals in the same Fungicide Group. The intrinsic risk for resistance to develop to a given fungicide group varies among chemistries; for example, resistance development among the strobilurins, Group 11, is much more likely than resistance development among the mancozeb or maneb, Group Y. For more information about Fungicide Resistance and the FRAC Fungicide list, see the following website: http://www.frac.info/frac_body.html.

Section 18 Emergency Uses Requested For 2009

See the ND Dept of Ag website for Section 18 (Emergency use) fungicides available in 2009.
<http://www.agdepartment.com/Programs/Plant/Section18Exemptions.html>

Fungicide Group Names

1. Methyl benzimidazole carbamates (MBC)
2. Dicarboximides
3. Demethylation inhibitors (DMI) (SBI: Class I)
4. Phenylamides (PA)
5. Amines (SBI: Class II)
6. Phosphoro-thiolates ; Dithiolanes
7. Carboxamides
8. Hydroxy-(2-amino-)pyrimidines
9. Anilino-pyrimidines (AP)
10. N-phenyl carbamates
11. Quinone outside inhibitors (QoI)
12. Phenylpyrroles (PP)
13. Quinolines
14. Aromatic hydrocarbons (AH); Heteroaromatics
15. Cinnamic acids
- 16.1. Melanin biosynthesis inhibitors-reductase (MBI-R)
- 16.2. Melanin biosynthesis inhibitors-dehydratase (MBI-D)
17. Hydroxyanilides (SBI: Class III)
18. SBI: Class IV
19. Polyoxins
20. Phenylureas
21. Quinone inside inhibitors (Qil)
22. Benzamides
23. Enopyranuronic acid antibiotic
24. Hexopyranosyl antibiotic
25. Glucopyranosyl antibiotic
26. Glucopyranosyl antibiotic
27. Cyanoacetamide-oximes
28. Carbamates
29. Unnamed
30. Organo tin compounds
31. Carboxylic acids
32. Heteroaromatics
33. Phosphonates
34. Phthalamic acids
35. Benzotriazines
36. Benzene-sulfonamides
37. Pyridazinones
- P. Several (host plant defense induction)
- U. Several
- M. Several (multi-site contact activity)