SEED TREATMENT

Chemical	Application	Dosage ¹	Control ² of Seedling Blight ³	Remarks
Captan Captan 400, 37.4%	Slurry	2-3.75 fl oz/cwt	Х	
Fludioxonil Maxim 4FS, 40.3%	Slurry	0.08-0.16 fl oz/cwt	X	
Mancozeb Dithane DF Rainshield NT, 75%	Slurry	2.1-4.3 oz/bu	X	
Dithane F-45, 37%	Drill box or slurry	3.2-6.4 fl oz/bu	X	
Dithane WSP or Penncozeb 80 WP, 80%	Drill box or slurry	2-4 oz/bu	×	
Penncozeb 75 DF, 75%	Drill box or slurry	2.1-4.3 oz/bu	×	
Manzate 75 ProStick, 75%	Slurry	2-4 oz/bu	X	
Thiram 42-S Thiram, 42%	Liquid or slurry	3 fl oz/bu	X	

Flax **FOLIAR SPRAYS**

Chemical (Fungicide Group)	Application ¹	Dosage ²	Pasmo (Septoria linicola) Control ³	Remarks
Pyraclostrobin (11) Headline, 23.6%	Spray	6-12 fl oz/A	Х	Resistance statement 5 ⁴ Apply at mid-flowering (7-10 days after flower initiation). Make 2 nd application of disease persists.

¹Dosage = Amount of formulated product to apply.

²X = Product labeled for crop and disease; Blank = product not labeled for specific disease.

³Seedling blights due to various fungal infections of seed.

Spray = ground or aerial; Fungigation = application through sprinkler irrigation system.

2Dosage = Amount of formulated product to apply.

3X = Product labeled for crop and disease; Blank = product not labeled for specific disease.

4Leaf spots includes fungal leaf diseases such as tan spot, Septoria blotch and spot blotch on wheat, and spot blotch and net blot on barley.

5See fungicide resistance management statements on pages 9 and 10.