



Seniors and Food Safety: To Market, To Market

Preventing Foodborne Illness

Prevention of foodborne illness starts with your trip to the supermarket. Pick up your packaged and canned foods first.

Don't buy food in cans that are bulging or dented or jars that are cracked or have loose or bulging lids. Look for expiration dates on the labels and never buy outdated food.

Likewise, check the "use by" or "sell by" dates on dairy products, such as cottage cheese, cream cheese, yogurt and sour cream, and pick the ones that will stay fresh longest in your refrigerator.

Choose eggs that are refrigerated in the store. Before putting them in your cart, open the carton to make sure none are cracked or leaking.

Save frozen foods and perishables such as meat, poultry or fish until last. Always put these products in separate plastic bags so that drippings don't contaminate other foods in your shopping cart.

Check for cleanliness at the meat or fish counter and the salad bar. For instance, cooked shrimp lying on the same bed of ice as raw fish could be contaminated. When buying from a salad bar, avoid fruits and vegetables that look brownish, slimy or dried out. These are signs that the product has been held at an improper temperature.

When shopping for shellfish, buy from markets that get their supplies from state-approved sources; stay clear of vendors who sell shellfish from roadside stands or the back of a truck. And if you're planning to harvest your own shellfish, heed posted warnings about the safety of the water.

Take an ice chest along to keep frozen and perishable foods cold if getting your groceries home will take more than two hours.



NDSU
Extension Service

North Dakota State University
 Fargo, North Dakota 58105

Safe Handling of Fruits and Vegetables

- Wash hands with warm water and soap for at least 20 seconds before and after handling food, especially fresh, whole fruits and vegetables, and raw meat, poultry and fish. Clean under fingernails, too.
- Rinse raw products in warm water. Don't use soap or other detergents. If necessary and appropriate, use a small scrub brush to remove surface dirt.
- Use smooth, durable and nonabsorbent cutting boards that can be cleaned and sanitized easily.
- Wash cutting boards with hot water, soap and a scrub brush to remove food particles. Then sanitize the boards by putting them through the automatic dishwasher or rinsing them in a solution of 1 teaspoon of chlorine bleach to 1 quart of water. Always wash boards and knives after cutting raw meat, poultry or seafood and before cutting another food to prevent cross-contamination.
- Store cut, peeled and broken-apart fruits and vegetables (such as melon balls) at or below 41 F in the refrigerator.

Julie Garden-Robinson, Ph.D., L.R.D.
Food and Nutrition Specialist

Source: Adapted with permission from FDA/ Center for Food Safety and Applied Nutrition, May 1999; www.cfsan.fda.gov/~dms/seniorsd.html
Developed in cooperation with AARP
This material is based upon work supported by the Cooperative State Research, Education and Extension Service, U.S. Department of Agriculture, under Agreement No. 2002-51110-01512.

Apply the Heat and Fight BAC!

Cooking food to the proper temperature kills harmful bacteria. This includes raw meat, poultry, fish and eggs, as well as foods that are thoroughly cooked upon purchase. Precooked foods may become contaminated during storage or handling.

Reheat ready-to-eat foods, such as hot dogs, luncheon meats, cold cuts, fermented and dry sausage, and other deli-style meat and poultry products, until they are steaming hot. If you cannot reheat these foods, do not eat them.

Cooking food to the proper temperature kills harmful bacteria. So Fight BAC!™ by thoroughly cooking your food as follows:

Raw Food	Internal Temperature
Ground Products	
Hamburger	160 F
Beef, veal, lamb, pork	160 F
Chicken, turkey	165 F
Beef, Veal, Lamb	
Roasts and steaks	
- <i>medium-rare</i>	145 F
- <i>medium</i>	160 F
- <i>well-done</i>	170 F
Pork	
Chops, roasts, ribs	
- <i>medium</i>	160 F
- <i>well-done</i>	170 F
Ham, fresh	160 F
Sausage, fresh	160 F
Poultry	
Chicken, whole and pieces	165 F
Duck	165 F
Turkey (unstuffed)	165 F
- <i>Whole</i>	165 F
- <i>Breast</i>	165 F
- <i>Dark meat</i>	165 F
- <i>Stuffing (cooked separately)</i>	165 F
Eggs	
Fried, poached	Yolk and white are firm
Casseroles	160 F
Sauces, custards	160 F

This chart has been adapted for home use and is consistent with consumer guidelines from the U.S. Department of Agriculture and U.S. Food and Drug Administration (FDA).

For more information about food safety, visit the NDSU Extension Service Web site: www.ag.ndsu.nodak.edu/food.htm

County commissions, North Dakota State University and U.S. Department of Agriculture cooperating. Duane Hauck, director, Fargo, N.D. Distributed in furtherance of the acts of Congress of May 8 and June 30, 1914. We offer our programs and facilities to all people regardless of race, color, national origin, religion, gender, disability, age, veteran's status or sexual orientation; and are an equal opportunity institution. This publication will be made available in alternative formats for people with disabilities upon request, (701) 231-7881.