

Keep Hot Foods Hot and Cold Foods Cold:


A Foodservice Guide to Thermometers and Safe Temperatures

Bacteria are found nearly everywhere, including in most foods. Bacteria double in number every 10 to 30 minutes when exposed to ideal temperatures, pH, nutrients or moisture levels. Controlling temperature is a way to prevent bacteria from growing and possibly causing food-borne illness.

Chefs should not depend on their instincts, cooking time, oven temperature or product appearance to determine when a product is done: thermometers are important tools for protecting foods.

Several types of thermometers are available for use with foods. Choose thermometers approved by the National Sanitation Foundation (NSF).

Glass mercury-filled thermometers should **never** be used for taking food temperatures because they could break. These thermometers should only be used for taking temperatures of people.



Types of thermometers

1. **Bimetal stemmed (or bayonet-style) thermometers** have a dial scale with a plastic lens cover and a temperature range from 0 to 220 degrees Fahrenheit (-18 to 105 C).
 - Their stems should be at least 5 inches long.
 - They should have a calibration nut under the dial.
 - They can accurately measure the temperature of relatively thick or deep foods (more than 2 inches) such as beef roasts and foods in stock pots.
2. **Refrigerator-freezer thermometers** may be built-in, hanging or sitting types.
 - They should be checked for accuracy using a calibrated stemmed thermometer: place the thermometers side by side in the refrigerator or freezer and compare readings. Inaccurate refrigerator-freezer thermometers usually cannot be recalibrated and must be replaced.
3. **Digital thermometers** show temperatures quickly on an easy-to-read display.
 - They are available in several sizes and styles, some with interchangeable temperature probes.
 - Some models can be calibrated. Change batteries based on use or according to indicator light.
4. **Candy, meat, microwave and deep-fry thermometers** are, as their names suggest, designed for special uses.

Foodservice/Restaurant Guide to Thermometers and Safe Temperatures

Calibrate stemmed thermometers regularly

Thermometers can get out of adjustment if they are jarred. Any time thermometers are dropped or exposed to extremes in temperatures, they should be calibrated.

All thermometers—regardless of use—should be calibrated at least once a month to maintain their accuracy.

Two acceptable methods for calibration are the Ice Point Method, which is recommended for high altitudes, and the Boiling Point Method.

Ice Point Method

1. Make a thick ice slush using crushed ice and cold water (50/50). Remove thermometer from its case and insert sensing area of thermometer into the center of the ice slush to a depth of at least two inches, avoiding the sides and bottom of the container.
2. Allow indicator to stabilize.
3. Adjust calibration nut so indicator reads 32 F (0 C).

Boiling Point Method

1. Insert sensing area of thermometer into boiling water avoiding sides and bottom of pan.
2. Allow indicator to stabilize.
3. Adjust calibration nut so indicator reads 212 F (100 C). To adjust for high altitudes, decrease setting 1 F (0.6 C) for each 550 feet above sea level.

Revised August 2007

This material was originally developed based upon work supported by the Extension Service, U.S. Department of Agriculture, under special project number 95-EFSF-0-3400.

For more information, visit www.ag.ndsu.nodak.edu/food.htm

County commissions, North Dakota State University and U.S. Department of Agriculture cooperating. Duane Hauck, director, Fargo, N.D. Distributed in furtherance of the acts of Congress of May 8 and June 30, 1914. We offer our programs and facilities to all people regardless of race, color, national origin, religion, gender, disability, age, veteran's status or sexual orientation; and are an equal opportunity institution. This publication will be made available in alternative formats for people with disabilities upon request, (701) 231-7881.

Care and Use of Thermometers

- Remove the protective case on stemmed thermometers.
- Wash thermometers in hot soapy water, rinse with clean hot water, sanitize with a solution of one tablespoon chlorine bleach per gallon of clean lukewarm (at least 75 F) water, and air-dry **before, between** and **after** uses. Keep the protective case clean, especially inside the case.
- Place thermometer in the center of the dish or in the thickest part of meat away from bone, inserting to the "dimple" or a minimum of two inches on the stem of the thermometer.
- Take the temperature reading when the needle on the dial has stopped moving for 15 seconds or the digital scale stabilizes.
- Take the temperature of the food in several places to confirm thorough cooking.
- Do not leave thermometers with plastic lens covers in foods during cooking. Use a meat thermometer or other thermometer that can withstand the heat of the oven.

Store, Cook and Hold Foods at Safe Temperatures

The Danger Zone refers to the temperature range from 41 to 135 degrees Fahrenheit, which provides ideal growth conditions for bacteria.

Cool foods from 140 to 70 F in less than 2 hours, and from 70 to 41 F within 4 hours for a total of 6 hours cooling time.

Safe Storage Temperatures

Coolers (Refrigerators)	41 F or below
Freezers	0 F or below

Safe Internal Cooking Temperatures

Ground beef, pork	155 F (for 15 seconds)
Poultry, stuffed meat, stuffed pasta, stuffing containing meat or poultry	165 F (for 15 seconds)
Beef Roasts	145 F (for 15 seconds)
	130 F (for 112 minutes)
Commercially processed, ready-to-eat food (cheese sticks)	135 F (for 15 seconds)
Cooked products to be reheated	165 F (for 15 seconds) (within 2 hours)
Eggs (immediate service)	145 F (for 15 seconds)
Eggs (hot held)	155 F (for 15 seconds)

* Measure the temperature of microwave-cooked foods in several places. Do not leave microwave thermometers in the food during cooking unless recommended by the manufacturer. Stir and rotate foods during cooking and allow to stand two minutes after cooking to allow for even distribution of heat.

Reference: FDA Food Code. 2005.

Prepared by **Julie Garden-Robinson**, Ph.D., LRD, Food and Nutrition Specialist